

LifeAssays® Equine SAA Technical Data:

- Easy-to-use
- Quantitative: clear relation of results vs. cut-off: 20 mg/L
- Accurate
- Rapid : assay time 5 minutes
- Small sample volume: 5 µL
- Linear range 10–1000 mg/L

LifeAssays® Magnia Reader Technical Data:

- Easy-to-read display
- Bench-top instrument
- Calibration keycard
- Automatic measurement and result presentation
- Superior non optical system
- Battery powered



References

1. Pepys MB, Baltz ML, Tennent GA, Kent J, Ousey J, Rossdale PD. Serum amyloid A protein (SAA) in horses: objective measurement of the acute phase response. *Equine Vet J.* 1989;21(2):106–109.
2. Hulten C, Tulamo RM, Suominen MM, Burvall K, Marhaug G, Forsberg M. A non-competitive chemiluminescence enzyme immunoassay for the equine acute phase protein serum amyloid A (SAA) - a clinically useful inflammatory marker in the horse. *Vet Immunol Immunopathol.* 1999;68(2-4):267–281. doi: 10.1016/S0165-2427(99)00027-6.
3. Cray C, Zaias J, Altman NH et al *Comp Med.* 59(6): 517–526, 2009 Dec ;. Acute Phase Response in Animals: A Review.
4. Jacobsen S. AAEP Proceedings, vol 53, 2007 Review of Equine Acute-Phase Proteins. <https://aaep.org/sites/default/files/issues/proceedings-07/proceedings-z9100107000230.pdf>
5. Tape C, Kisilevsky R. Apolipoprotein A-I and apolipoprotein SAA half-lives during acute inflammation and amyloidogenesis. *Biochim Biophys Acta.* 1990;1043(3):295–300.
6. Jacobsen S, Nielsen JV, Kjelgaard-Hansen M, Toelboell T, Fjeldborg J, Halling-Thomsen M, Martinussen T, Thoenfer MB. Acute phase response to surgery of varying intensity in horses: a preliminary study. *Vet Surg.* 2009;38(6):762–769. doi: 10.1111/j.1532-950X.2009.00564.x
7. Hobo S, Niwa H, Anzai T. Evaluation of serum amyloid A and surfactant protein D in sera for identification of the clinical condition of horses with bacterial pneumonia. *J Vet Med Sci.* 2007;69(8):827–830. doi: 10.1292/jvms.69.827
8. Jacobsen S, Thomsen MH, Nanni S. Concentrations of serum amyloid A in serum and synovial fluid from healthy horses and horses with joint disease. *Am J Vet Res.* 2006;67(10):1738–1742. doi: 10.2460/ajvr.67.10.1738.
9. Paltrinieri S, Giordano A, Villani M, Manfrin M, Panzani S, Veronesi MC. Influence of age and foaling on plasma protein electrophoresis and serum amyloid A and their possible role as markers of equine neonatal septicemia. *Vet J.* 2008;176(3):393–396. doi: 10.1016/j.tvjl.2007.05.018.
10. Hulten C, Sandgren B, Skioldebrand E, Klingeborn B, Marhaug G, Forsberg M. The acute phase protein serum amyloid A (SAA) as an inflammatory marker in equine influenza virus infection. *Acta Vet Scand.* 1999;40(4):323–333.